

AD-A056 844

CALIFORNIA UNIV LOS ANGELES PLASMA PHYSICS GROUP
EXCITATION OF LOWER HYBRID WAVES IN A FINITE PLASMA.(U)
FEB 78 V K DECYK, J M DAWSON, G J MORALES

F/G 20/9













UNCLASSIFIED

PPG-338

N00014-75-C-0476

NL



1 OF 1
ADA
056844



END
DATE
FILMED
9 -78
DDC

AD No. _____
DDC FILE COPY

AD A 056844



CENTER FOR
PLASMA PHYSICS
AND
FUSION ENGINEERING

UNIVERSITY OF CALIFORNIA
LOS ANGELES

12

LEVEL II

DDC
RECEIVED
JUL 31 1978
B

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

21 066

AD A 056844

12



LEVEL II

6

EXCITATION OF LOWER HYBRID WAVES
IN A FINITE PLASMA

9

Doctoral thesis,

10

V. K./Decyk, J. M./Dawson, G. J./Morales

14

PPG-338

11

Feb 1978

12

44 p.

15

N00014-75-C-0476, EY-76-C-03-0010

AD No.

DC FILE COPY

Department of Physics
University of California
Los Angeles, California 90024

DDC
RECEIVED
JUL 31 1978
B

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

78 07 21 066

401733

act

ABSTRACT

A computer simulation study of the launching of lower hybrid waves by external sources and their propagation in a finite but homogeneous plasma slab is presented. The study makes use of a 2-1/2 dimensional, electrostatic particle code developed for simulation of bounded plasma. A number of antenna configurations are considered: a point source, two finite-length capacitor plates, and a phased array of capacitor plates. When the oscillating frequency does not match a bounded plasma resonance, one can observe resonance cones, energy absorption at the plasma surface, ion cyclotron modulation of the source, and energetic ions, depending on the parameters chosen. The excitation of a bounded plasma resonance is also considered. The nonlinear evolution of the resonance shows that wave-particle interactions and ponderomotive force effects play an important role. The possibility of controlling the electron temperature profile is discussed.

ACCESSION for	
NTIS	Write Section <input checked="" type="checkbox"/>
DDC	Self Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist. and/or SPECIAL	
A	-

I. INTRODUCTION

Plasma heating by lower hybrid waves has been considered extensively over the past decade.¹⁻¹⁴ The technological advantages of this scheme are: (1) RF generators exist which can deliver large amounts of power at the frequencies required; (2) the coupling to fusion devices can be implemented through waveguides which are remote from the neutron environment. However, if this scheme is to be useful, a number of plasma physics questions must be solved. There is the problem of effectively coupling the RF energy to the plasma at the edge, especially at the power levels required.^{2,3} Further, one must understand what happens to the wave as it propagates to the resonance layer, especially at large wave amplitudes, where many nonlinear effects may be important.⁴⁻⁷ Thirdly, if the RF energy can penetrate to the resonance region, there is the problem of what linear and/or nonlinear effects occur in the lower hybrid regions.^{1,8-12} Finally, the processes by which the energy of the wave is absorbed by the plasma, and the subsequent thermalization of the energy must be understood.¹¹⁻¹³

Computer simulation can be a useful tool in understanding the physics in this problem, especially when nonlinear effects are important and are difficult to treat analytically. Present day computers are not large enough nor fast enough, however, to handle all the different scale lengths necessary in simulating the entire lower hybrid heating scheme. Instead, one must at present be satisfied with simulating certain parts of the scheme separately, such as the coupling of the radiation at the edge, or the wave conversion in the plasma interior. One can nonetheless identify the important kinetic and nonlinear effects in each area and thus contribute to the understanding of the overall problem. As a first step in a systematic study of the lower hybrid heating scheme, we present here a

computer simulation study of the launching and propagation of lower hybrid waves by external sources and their propagation in a finite but homogeneous plasma slab bounded on both sides by vacuum.

The computer model and simulation parameters are discussed in Sec. II. Non-resonant oscillations excited by various sources are discussed in Sec. III. In Sec. IV the excitation and nonlinear evolution of a bounded plasma resonance are studied. Finally, a summary is contained in Sec. V.

II. COMPUTER MODEL

An electrostatic particle simulation model, described in detail elsewhere,¹⁵ has been developed for this study and is briefly summarized here. The two dimensional electrostatic model is periodic in one dimension and bounded in the other. In the non-periodic dimension one can specify the values of the potential or the normal electric field on both boundaries. It is also possible to specify that a vacuum exists outside the plasma, or any combination of these conditions on each boundary separately. External electrostatic sources (i.e. charges not belonging to the plasma) can be added in a natural way either on the boundaries or in the plasma interior. Finite-sized (Gaussian) particles are used throughout, and all three velocity co-ordinates are retained in the calculations.

For the simulation results presented here, the following parameters and conditions were specified: Vacuum boundary conditions were imposed in the \hat{x} direction on both sides of the slab. External sources, described below, are specified on one boundary. The external magnetic field was oriented in the periodic (\hat{z}) direction with ratio of electron cyclotron to plasma frequency $\Omega_e/\omega_{pe} = 1$, and ion-electron mass ratio $M_i/m_e = 64$. In most cases the ions were colder than electrons with $T_e/T_i = 10$ (except for the case of Fig. 4 where $T_e = T_i$). Particles were initially uniformly distributed within the slab with Gaussian velocity distributions, and they were reflected elastically from the boundaries, with the full dynamics calculated correctly to second order in the time step $\Delta t = 0.2 \omega_{pe}^{-1}$. The reflection of particles implies that there is no mechanism for net energy loss in the system. The size of the system in the \hat{x} and \hat{z} directions was $L_x/\lambda_{De} = 32$, $L_z/\lambda_{De} = 128$.

The external source in these simulations was determined by specifying an external surface charge density on one boundary of the slab, oscillating

with a specified frequency ω_0 . Because the magnitude of the charge density is fixed externally, such a source corresponds to a constant current driver. In all cases, the phase of the oscillator was chosen so that the external field was zero at $t = 0$, and the amplitude was kept small so that the plasma response could be described, at least initially, by a linear theory. The strength of the oscillator was normalized in terms of a parameter w , the energy of the source in vacuum integrated over the plasma area A divided by the electric field fluctuation energy of the plasma when in thermal equilibrium and free of external sources,

$$w \equiv \frac{1}{A} \int E_{\text{ext}}^2 dA / \sum_k |E_k|^2. \quad (1)$$

In the computer model for the parameters chosen, the thermal electric field fluctuation energy density is about 100 times smaller than the particle kinetic energy density. The reason for this particular normalization is that the plasma response is determined by the length and distribution of the external charge, as well as its magnitude, and all these factors are included in w .

III. NON-RESONANT OSCILLATIONS DRIVEN BY AN EXTERNAL SOURCE

It is well-known that the normal modes of oscillation of a bounded plasma form a discrete set.¹⁶ Therefore, the plasma will respond differently depending on whether or not the frequency and a wavenumber of the spectrum excited by the source match simultaneously the frequency and wavenumber of one of the normal modes of the plasma. For perfect matching, the electric field for the single resonant mode grows secularly with time until some nonlinearity stops the growth. If there is a mismatch, the plasma responds with non-resonant driven oscillations. Since a finite size driver contains a spectrum of wavenumbers, there will generally be a response of many non-resonant modes even when one mode satisfies the resonance condition.

The typical non-resonant plasma response to a number of "antenna configurations" of external sources will be considered next, along with the significant kinetic and nonlinear effects. Special attention will be paid to the coupling of the source to the plasma at the surface, the spatial distribution of the heating, and the evolution of the velocity distribution.

A. Point Source Antenna and Lower Hybrid Resonance Cones

A well-known feature of the propagation of lower hybrid signals is the existence of resonance cones.^{4,5} These cones arise because for a given frequency, the group velocities away from the source for all wavenumbers with the same sign are parallel, in the cold plasma limit. In order to clearly observe resonance cones in a model which is periodic in one direction, it is necessary that the cones from each of the periodic sources must interfere constructively. In addition, the frequency and wavenumbers of the source should not correspond to some bounded plasma mode in order to avoid a single mode dominating the plasma response. Since a resonance cone is composed of the sum of many Fourier modes, a point source (actually a line source in

three dimensions) was chosen as the driver in this case, because it contains a broad band spectrum. A driving frequency which satisfies the above conditions was then chosen ($\omega_0 = 2.7 \omega_{pi}$). The parameter w specifying the strength of the driver was set equal to $2/3$.

The system was followed for about 11 oscillation periods. It was found that the resonance cones were set up in about one oscillation period, but the amplitude of the potential oscillations was rather weak. In order to see the cones clearly, the potential at every point in space was Fourier analyzed in time to extract the plasma response at the driver frequency ω_0 . The result is shown in Fig. 1. The two cones are clearly seen, and make an angle $\theta \approx 27^\circ$ with respect to the magnetic field; however, they are broader than expected from cold plasma theory. The reason the cones are broad is that the shorter wavelength components of the potential are quickly absorbed by electron Landau damping at the surface.

The resonance cones shown in Fig. 1 can be Fourier analyzed in the parallel direction. This analysis shows that although the point source has a broad spectrum of wavenumbers, only the three largest wavelength modes appreciably contribute to the cones in the plasma interior. The remaining modes, whose parallel phase velocities satisfy $\omega_0/k_m \lesssim 2.5 v_{th}$, are absorbed at the surface. (The parallel wavenumber is given by $k_m = 2m\pi/L_z$ for integer m and $v_{th} = \sqrt{kTe/m_e}$ is the electron thermal velocity.)

The simulation results reveal a number of other important effects. In Fig. 1, one can observe the dramatic potential drop due to the plasma sheath surrounding the source. The effect of the sheath is to further reduce the penetration of the source. Examination of the time evolution of the total electric field energy w_{ef} , shown in Fig. 2 and normalized to the kinetic energy of a thermal electron by

$$w_{ef} \equiv \frac{\tilde{\lambda}_{De}^2}{kT_e} \int \frac{E^2}{8\pi} dA, \quad (2)$$

where $\tilde{\lambda}_{De}$ is the dimensionless Debye length, reveals another interesting kinetic effect. Superimposed on the driven oscillations, there is a modulation of the pump at twice the ion cyclotron frequency Ω_i , which manifests itself as a modulation of the electric field energy at $4\Omega_i$. For all the non-resonant cases studied, a modulation at the ion cyclotron frequency or its first harmonic was observed whenever the physical length of the driver was comparable or smaller than the ion Larmor radius. Although this modulation effect needs further study, it is probably due to ion kinetic effects (ion Bernstein modes) being excited in the sheath region, perhaps due to the sudden turning on of the pump.

The kinetic energy of each species α is normalized in the same manner as the electric field energy, according to

$$w_\alpha \equiv \frac{\tilde{\lambda}_{De}^2}{kT_e} \frac{1}{2} m_\alpha \sum_j v_j^2 \quad (3)$$

where the sum is over the particles of species α . The evolution of the electron kinetic energy w_e (Fig. 3) shows a net energy gain of about 5% after 11 oscillation periods, with the gain all due to the increase of parallel velocity. The slight undulations in the growth coincide with the ion cyclotron harmonic modulation noted previously. The distribution of energy was not spatially uniform, however. The parallel thermal velocity of electrons confined to the region near the source increased by 20%, whereas the thermal velocity of particles on the opposite side of the slab increased less than 2%. Although a few fast electrons were formed, the heating occurred mainly in the bulk of the velocity distribution. The rate of energy gain of the cold ions, also illustrated in Fig. 3, was given by

$$(\omega_{pe} T_e)^{-1} \frac{dT_i}{dt} \approx 7 \times 10^{-5} . \quad (4)$$

This is about the same rate observed when no source is present. Further,* the energy of the ions stayed spatially uniform. This indicates that the energy gain of the ions must have been due to collisional heating with the warm electrons. The driving frequency in this case was too high for much direct ion response.

B. Short Capacitor Plates as Antenna

For the case of the point source, it is apparent that electron Landau damping at the surface plays an important role in the excitation of lower hybrid waves in the plasma interior. This "filtering" effect of the plasma surface is even more dramatically demonstrated here, where two short capacitor plates oscillating out of phase are used as the external source. The frequency ($\omega_o = 2\omega_{pi}$) and length of the source was chosen so that the wavenumber spectrum had a maximum around those modes whose parallel phase velocities were near the electron thermal velocity v_{th} , and the strength of the driver was chosen so that $w = 4/3$. The equipotential lines for the capacitor plates in vacuum are illustrated in Fig. 4(a).

The computer plasma was followed for about 12 oscillation periods, and the steady-state plasma response at the driving frequency is shown in Fig. 4(b). The simulation results show that most of the modes are absorbed at the plasma surface, and only the longest wavelength mode is able to penetrate to the plasma interior. Figure 4(c), which shows the plasma response predicted by cold plasma theory for the excitation of the single mode $m = 1$, verifies that indeed only that mode was excited in the simulation. In other respects, such as the presence of the plasma sheath surrounding the exciter, the modulation of the source at the ion cyclotron frequency, and the evolution of the plasma heating, the results here are similar to those of the point source.

C. Phased Array of Capacitor Plates as Antenna

A phased array of capacitor plates has also been used as the exciting structure. The frequency ($\omega_o = 2\omega_{pi}$) and length of each element was chosen so that the wavenumber spectrum was sharply peaked around a single mode whose parallel phase velocity was $\omega_o/k_m = 1.3 v_{th}$, which should lead to a strong interaction of the electrons with the exciting structure. To enhance this interaction, the strength of the driver was increased to $w = 10$. The equipotential lines for the phased array in vacuum is illustrated in Fig. 5(a). The computer plasma was followed for 12 oscillation periods, and as shown in Fig. 5(b), there exist no driven oscillations in the plasma interior.

The major observations in this case are the familiar electron surface heating and the presence of significant ion heating, which is also largest at the plasma surface. The electron kinetic energy w_e , illustrated in Fig. 6, shows an energy gain of 20% after $\omega_{pe} t = 300$, mostly due to an increase in the parallel velocity. The energy gain was localized in space to within a few electron Larmor diameters of the source, as shown in Fig. 7.

The ion heating, on the other hand, shows a dramatic increase over previous levels. The ion kinetic energy w_i , also shown in Fig. 6, increases by nearly 200% after $\omega_{pe} t = 300$. Like the electron case, the kinetic energy distribution is inhomogeneous in space and decreases away from the source (Fig. 7). However, unlike the situation for electrons, the energy gain for the ions was mostly due to an increase in velocity in the \hat{x} direction, perpendicular to the magnetic field. Furthermore, as shown in Fig. 8, energetic ion tails are formed in the ion velocity distribution $f_i(v_x)$. The reason for this difference in ion behavior is that the large perpendicular electric fields near the exciting structure (Fig. 5(b)) strongly accelerate the ions but do not affect the magnetized electrons. The magnetic field would eventually return the ions back to the exciter and they could be accelerated again.

IV. RESONANT OSCILLATIONS DRIVEN BY AN EXTERNAL SOURCE

In the previous section, non-resonant driven oscillations were observed to have generally low amplitudes inside the plasma. In this section, the enhanced plasma response when the external source excites a bounded plasma resonance is considered. One example is studied in detail, and particular attention is paid to the nonlinear evolution of the system and the space and velocity-space distribution of the plasma heating.

The dispersion relation of the bounded plasma slab, based on a warm electron-cold ion fluid model was calculated¹⁷ and is illustrated in Fig. 9. Here the frequency is plotted as a function of the parallel wavenumber and each curve corresponds to different allowable perpendicular wavenumbers. The frequency ($\omega_o = 2.3 \omega_{pi}$) and length of two phased capacitor plates was chosen so that the wavenumber spectrum peaks at a normal mode (point A in Fig. 9) whose parallel phase velocity is about $6 v_{th}$, and thus there would be little electron heating initially by Landau damping. The source strength was chosen to be $w = 1$.

The computer plasma was followed for 27 oscillation periods. The time evolution of the electric field energy, illustrated in Fig. 10, shows the secular resonant growth followed by absorption of the wave energy. Then somewhat later, the process repeats. The external source, whose amplitude is indicated by the arrow in Fig. 10, was kept on during the entire run. Examination of the evolution of the total kinetic energy of each species, illustrated in Fig. 11, shows that the energy absorption is dominated by electrons.

A. Electron Dynamics

The rate of increase of the electron kinetic energy and the amplitude of the electric field energy are found to be closely correlated, both reaching their maximum value at $\omega_{pe} t = 85$ and their minimum value at

$\omega_{pe} t = 130$. By examining the electron velocity distribution function, one finds that the increase in kinetic energy is entirely due to electrons quickly being accelerated to large velocities along the \hat{z} direction, as shown in Fig. 12. One can see that electrons having velocities as large as $10 v_{th}$ (that is, with 100 times the thermal energy) are created by $\omega_{pet} = 220$. This process continues until the resonant wave has lost most of its energy.

This behavior can be understood in terms of the concept of particle trapping. Initially there were no electrons moving close enough to the phase velocity of the wave to be trapped. As the potential ϕ_0 grows in magnitude, a few particles can satisfy the trapping criterion

$$v > \frac{\omega}{k} - 2\sqrt{\frac{e\phi_0}{m_e}} \quad (5)$$

When this happens, the wave accelerates the particles and loses some energy. If the wave damping is small, the wave will regain this energy when the trapped particles are later decelerated by the wave. However, if many particles are suddenly accelerated, the wave may lose so much energy that it can no longer trap the particles, and the result is the generation of fast particles accompanied by the decay of the wave. Tail formation was observed to occur in the simulation when

$$\frac{\omega}{k} - 2\sqrt{\frac{e\phi_0}{m_e}} \approx 2.5 v_{th} \quad (6)$$

indicating that this acceleration process occurred when the wave amplitude was large enough to begin to trap particles in the bulk of the electron velocity distribution. A similar mechanism has been discussed by Dawson and Shanny.¹⁸

A significant feature of the interaction of the electrons with the wave is that the interaction is localized in space because the eigenmode (not shown) has large parallel electric fields near the boundaries. Since the electrons are tightly bound to the magnetic field lines, it is not the spatial distribution of the total electric field for the eigenmode but rather the distribution of the parallel electric field which is important in accelerating the electrons, even though the magnitude of the parallel electric field is small compared to the perpendicular electric field. A graph of the electron kinetic energy versus position in the perpendicular direction, Fig. 13, shows that the kinetic energy profile in space follows approximately the same pattern as the parallel electric field energy for the eigenmode (not shown).

These results illustrate a possible method for controlling the electron temperature profile in a bounded plasma by exciting eigenmodes whose wavelengths are comparable to the plasma width. Specifically, since a bounded plasma mode is determined by the plasma parameters and geometry, one can obtain a desired temperature profile if a mode with appropriate parallel electric field distribution is chosen. For the case illustrated in Fig. 13, about equal electron heating on both sides of the plasma was obtained, even though the exciter was only on one boundary. Of course, only the parallel energy is increased.

B. Ion Dynamics

The ions do not gain much energy from the resonant wave, as one can see from Fig. 11. The phase velocity of the excited wave is too fast to resonate directly with the ions, and except for the increase of oscillation energy evident in Fig. 11, the ions can only gain energy in a nonlinear fashion. The key to understanding the behavior of the ions turns out to be the ponderomotive force on the electrons and the resulting density depressions. Furthermore, by

understanding the ion behavior the reason for the second increase in wave amplitude, shown in Fig. 10, becomes clear.

Since the normal mode being excited has a strong spatial distribution of electric field energy, the electrons feel a large ponderomotive force. The largest component for electrons is due to the parallel gradients of the parallel electric field energy.⁶ For ions the ponderomotive force is negligibly small. The electrons are therefore pushed away from regions of large parallel electric field, and the resulting charge imbalance gives rise to an ambi-polar potential which forces electrons and ions out of the high field regions together, leaving a density depression.

Ions streaming away from the high field regions are observed in the simulation beginning at $\omega_{pe} t = 100$, and there is a corresponding increase in ion kinetic energy (Fig. 11). The peak density depression occurs around $\omega_{pe} t = 240$, and a three-dimensional view of the density (for electrons) is shown in Fig. 14 (averaged over one oscillation period). The ion density looks similar, although somewhat noisier. This mechanism of ion heating has also been observed in recent experiments investigating the focus region of intense RF radiation above the lower hybrid frequency.¹⁹ The electrons gain a negligible amount of energy in this manner compared with the energy they gain directly from the wave.

Because the system is periodic in the \hat{z} direction, ions from adjacent high field regions accumulate between these regions, until a potential develops that opposes further ion build-up. Shortly after, around $\omega_{pe} t = 240$, the ion kinetic energy shows another jump because even though some ions are slowed by this potential, others are accelerated even more. Thus a slowly-varying potential modulates the ion heating, and this shows up as undulations in Fig. 11. This type of ion heating resulted in fast tails in the parallel velocity distribution for ions, with ions having velocities as large as 5 or 6 times the background ion thermal velocity.

C. Relaxation Oscillations

When the plasma density relaxes to a nearly homogeneous state (although fluctuations have increased), around $\omega_{pe} t = 350$, the electric field returns to resonant growth. This recurrence can be described by the term relaxation oscillation. Although the external source has been on all this time, the plasma density had changed so much that the source was no longer coupling to a resonant mode. But when the density finally relaxes, a resonant mode can be excited again. The whole process we have been describing repeats itself, although the growth this time is not so rapid, since there are now particles traveling fast enough to interact with the wave.

Although in the example we have been studying, it has been the sudden acceleration of electrons which damped the resonant wave, it is conceivable that with different parameters (such as a resonant mode with a higher parallel phase velocity), the density modification could have "detuned" the resonance first, in which case the evolution of the plasma heating could have been quite different. This possibility needs further study, because it might lead to larger ion heating.

The plasma state at the end of the computer run has several times larger fluctuations of density and electric field than thermal level. Frequency analysis indicates that parametric decay to other modes was not significant; for the two dimensional geometry chosen, however, such decay was not expected.

V. SUMMARY

Summarizing the results for the non-resonant driven oscillations at frequencies above the lower hybrid, the plasma response in the interior is relatively small. At the surface, the plasma response is dominated by electrons. Resonance cones can be excited in this model at certain frequencies. In all cases a large potential drop near the exciter was observed due to the presence of a plasma sheath. Modulation of the source at ion cyclotron harmonics by the plasma was also observed. The plasma surface acts as a filter, absorbing those wavenumbers whose phase velocities fall in the electron velocity distribution. Most of the plasma heating occurs at the surface and takes the form of an increase of electron velocity parallel to the magnetic field but without the creation of fast tails in the velocity distribution. Ions were appreciably heated only for relatively large amplitude exciters by being accelerated away from the source.

The results for the resonant driven oscillations can be described as initial growth followed by relaxation oscillations. The evolution of the energy is dominated by electrons. The resonance is quenched by sudden tail formation and resultant wave damping. Electron heating is localized in space and follows the pattern of the parallel electric field energy for the resonant mode. This offers a possibility for controlling the electron temperature profile in space by excitation of bounded plasma resonances. Nonlinear density changes associated with the ponderomotive force are significant. Ions gain energy by being expelled from the density cavities by the ambi-polar potential associated with the ponderomotive force, as observed in recent experiments. When the density relaxes, the resonant mode is excited again, and the process repeats.

ACKNOWLEDGMENTS

We wish to thank Dr. William Kruer for useful discussions concerning the simulation results. The contents of this paper form part of a dissertation submitted by one of the authors (V. K. D.) to the University of California, Los Angeles, in partial fulfillment of the requirements for the Ph. D. degree. This work is supported by U. S. D. O. E. Contract No. EY-76-C-03-0010 PA26/TASK III, and ONR Contract No. N00014-75-C-0476 P00004.

REFERENCES

1. T. H. Stix, Phys. Rev. Lett. 15, 878 (1965).
2. V. E. Golant, Zh. Tekh. Fiz. 41, 2492 (1971) [Sov. Phys.-Tech. Phys. 16, 1980 (1972)].
3. G. J. Morales, Phys. Fluids 20, 1164 (1977).
4. R. F. Fisher and R. W. Gould, Phys. Fluids 14, 857 (1971).
5. P. M. Bellan and M. Porkolab, Phys. Fluids 19, 995 (1976).
6. G. J. Morales and Y. C. Lee, Phys. Rev. Lett. 35, 930 (1975).
7. R. L. Stenzel and W. Gekeleman, Phys. Fluids 20, 108 (1977).
8. V. E. Golant and A. D. Piliya, Usp. Fiz. Nauk 104, 413 (1971) [Sov. Phys.-Usp. 14, 413 (1972)].
9. J. M. Kindel, H. Okuda, and J.M. Dawson, Phys. Rev. Lett. 29, 995 (1972).
10. M. Porkolab, Phys. Fluids 17, 1432 (1974).
11. A. Hasegawa and L. Chen, Phys. Fluids 18, 1321 (1975).
12. C. Chu and J. M. Dawson, Phys. Fluids 19, 981 (1976).
13. M. Brambilla, Plasma Phys. 18, 669 (1976).
14. M. Porkolab, S. Bernabei, W. M. Hooke, R. W. Motley, and T. Nagashima, Phys. Rev. Lett. 38, 230 (1977).
15. V. K. Decyk and J. M. Dawson, UCLA Center for Plasma Physics and Fusion Engineering Report PPG-331 (1977). To be published.
16. C. C. Cheng and E. G. Harris, Phys. Fluids 12, 1262 (1969).
17. Viktor K. Decyk, Ph. D. dissertation, UCLA (1977).
18. J. M. Dawson and R. Shanny, Phys. Fluids 11, 1506 (1968).
19. W. Gekeleman and R. L. Stenzel, Phys. Fluids 20, 1316 (1977).

FIGURES

- FIG. 1 Electrostatic potential associated with lower hybrid resonance cones excited by point source. ($\Omega_e/\omega_{pe} = 1$, $M_i/m_e = 64$, $T_e/T_i = 10$, $\omega_o/\omega_{pe} = .35$, $w = 2/3$, $2 \times 42 \times 192$ particles, 32×128 grid, particle size $a_x = a_z = 1$, $\tilde{\lambda}_{De} = 1$)
- FIG. 2 Time evolution of the total electric field energy for a point source exciter.
- FIG. 3 Time evolution of the scaled electron (w_e) and ion (w_i) kinetic energies for a point source exciter.
- FIG. 4 Equipotential lines for:
- (a) short capacitor plates in vacuum;
 - (b) oscillations driven by short capacitor plates as observed in simulation; ($\Omega_e/\omega_{pe} = 1$, $M_i/m_e = 64$, $T_e/T_i = 1$, $\omega_o/\omega_{pe} = .25$, $w = 4/3$, $2 \times 42 \times 192$ particles, 32×128 grid, particle size $a_x = a_z = 1$, $\tilde{\lambda}_{De} = 1$);
 - (c) oscillations driven by short capacitor plates as predicted by fluid model.
- FIG. 5 Equipotential lines for:
- (a) phased array of capacitor plates in vacuum;
 - (b) oscillations driven by phased array of capacitor plates as observed in simulation; ($\Omega_e/\omega_{pe} = 1$, $M_i/m_e = 64$, $T_e/T_i = 10$, $\omega_o/\omega_{pe} = .25$, $w = 10$, $2 \times 42 \times 192$ particles, 32×128 grid, particle size $a_x = a_z = 1$, $\tilde{\lambda}_{De} = 1$).

- FIG. 6 Time evolution of scaled electron (w_e) and ion (w_i) kinetic energies for a phased array of capacitor plates.
- FIG. 7 Spatial dependence of the scaled electron (w_e) and ion (w_i) kinetic energies perpendicular to the magnetic field (source is at $x = 0$).
- FIG. 8 Ion velocity distribution perpendicular to the magnetic field, averaged over all space. Dashed curve corresponds to initial distribution and solid curve to final distribution.
- FIG. 9 Calculated dispersion relation for a plasma slab with cold ions and $L_x/\lambda_{De} = 32$.
- FIG. 10 Time evolution of the electric field energy for the resonantly driven case. Arrow indicates energy of exciter. ($\Omega_e/\omega_{pe} = 1$, $M_i/m_e = 64$, $T_e/T_i = 10$, $\omega_o/\omega_{pe} = .28$, $w = 1$, $2 \times 42 \times 192$ particles, 32×128 grid, particle size $a_x = a_z = 1$, $\tilde{\lambda}_{De} = 1$)
- FIG. 11 Time evolution of the scaled electron (w_e) and ion (w_i) kinetic energies for the resonantly driven case.
- FIG. 12 Electron velocity distribution parallel to the magnetic field, averaged over all space. Dashed curve corresponds to initial distribution and solid curve to distribution after wave damping.
- FIG. 13 Spatial dependence of the electron kinetic energy perpendicular to the magnetic field. (Source is at $x = 0$.) Bottom curve shows energy dependence after first wave damping and top curve shows energy dependence after second wave damping.
- FIG. 14 Electron density profile at peak density depression, averaged over one oscillation period.




FIGURE 1




FIGURE 2




FIGURE 3






FIGURE 4





FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14

UCLA PLASMA PHYSICS GROUP REPORTS

- PPC-69 "Propagation of Ion Acoustic Waves Along Cylindrical Plasma Columns", *Phys. Fluids* **9**, 1281 (1966) (A.Y. Wong).
PPC-70 "Stability Limits for Longitudinal Waves in Ion Beam-Plasma Interaction", *Phys. Fluids* **9**, 1084 (1966) (B.D. Fried and A.Y. Wong).
PPC-71 "The Kinetic Equation for an Unstable Plasma in Parallel Electric and Magnetic Fields", *Phys. Fluids* **9**, 2428 (1966) (B.D. Fried and S.L. Osakow).
PPC-72 "Low-Frequency Spatial Response of a Collisional Electron Plasma", *Phys. Fluids* **9**, 292 (1966) (B.D. Fried, A.N. Kaufman and S.L. Osakow).
PPC-73 "Effects of Collisions on Electrostatic Ion Cyclotron Waves", *Phys. Lett.* **25**, 157 (1966) (A.Y. Wong, D. Judd and F. Hal).
PPC-74 "Interaction Between Ion Beams and Plasmas", *Bull. Am. Phys. Soc.*, **11**, 1182 (1966) (R. Roehrig, A.Y. Wong and J.M. Seiten).
PPC-75 "Excitation and Damping of Drift Waves", *Phys. Rev. Lett.* **21**, 157 (1966) (D.E. Kaplan and I.M. H).
PPC-76 "The Guiding Center Approximation in Lowest Order", *J. Plasma Phys.* **1**, 305 (1967) (A. Benio, Jr.).
PPC-77 "Plasma Streaming into a Magnetic Field", *J. Plasma Phys.* **1**, 305 (1967) (A. Benio, Jr.).
PPC-78 "Cooperative Effects in Plasma Echo Phenomena", *Proc. of 1st Int'l. Conf. on Quasilinear Plasmas*, Frascati, Rome, 1967, (A. Benio, Jr.).
PPC-79 "A Quasilinear Study of the Electron Gas Via the Test Particle Method", Dissertation, March 1967 (M.E. Rensink).
PPC-80 "Linear and Nonlinear Theory of Grid Excitation of Low Frequency Waves in a Plasma", Dissertation, April 1967 (G.L. Johnston).
PPC-81 "The Expansion and Diffusion of an Isolated Plasma Column", Dissertation, May 1967 (J. Hyman).
PPC-82 "Two-Pole Approximation for the Plasma Dispersion Function", *Phys. Fluids* **10**, 249 (1968) (B.D. Fried, C.L. Hedrick and J. McCune).
PPC-83 "Experimental Investigation of Electron Beam Phenomena", Dissertation, August 1967 (J.S. DeGroot).
PPC-84 "Cyclotron Echoes from Doppler Effects", *March 1968* (A.Y. Wong).
PPC-85 "Cyclotron Echoes from Doppler Effects", *March 1968* (A.Y. Wong).
PPC-86 "Ion Wave Echoes", *Phys. Rev. Lett.* **20**, 318 (1968) (D.R. Baker, R.R. Ahern and A.Y. Wong).
PPC-87 "Artificial Van Allen Belt", *Phys. Rev. Lett.* **21**, 528 (1968) (A.Y. Wong and R. Roehrig).
PPC-88 "Landau Damping of Ion Acoustic Waves in a Cesium Plasma with Variable Electron-Ion Temperature Ratio", Dissertation, October 1967 (G. Boer).
PPC-89 "The Two-Stream Instability", September 1967 (G. Boer).
PPC-90 "Magnetic Turbulence in Shocks", *Physics of the Magnetosphere*, R. Carovillano, J.F. McClay and H.R. Rudoski, eds. (G. Reidel, Dordrecht, Holland, 1969), pp. 485-513 (G.F. Kennel and H.E. Petschek).
PPC-91 "Small Amplitude Waves in High Beta Plasmas", *J. Plasma Phys.* **3**, 55 (1969) (O. Fornasiero and C.F. Kennel).
PPC-92 "Low Beta Plasma Penetration Across a Magnetic Field", *Phys. Fluids* **12**, 702 (1969) (B.D. Fried and S. Osakow).
PPC-93 "Annual Status Report", Feb. 1, 1967-Jun. 30, 1968, Principal Investigators, A. Mann, Jr., B.D. Fried, and C.F. Kennel.
PPC-94 "The Theoretical Investigation of Ionospheric Space", *Plasma Waves in Space and in the Laboratory*, ed. J.O. Thomas and B.J. Leubner (Edinburgh U. Press, Edinburgh, 1969), Vol. 11, (C.F. Kennel and F.L. Scarf).
PPC-95 "Magnetospheric Echoes in Collisionless Plasmas", *Phys. Fluids* **12**, 866 (1969) (A.Y. Wong).
PPC-96 "Parametric Excitation of Drift Waves in a Resistive Plasma", *Phys. Fluids* **12**, 1097 (1969) (G. Weyl and M. Goldman).
PPC-97 "Current Decay in a Streaming Plasma Due to Ion Turbulence", *Phys. Fluids* **12**, 274 (1969) (S.L. Osakow and B.D. Fried).
PPC-98 "Temperature Gradient Instabilities in Anisotropic Systems", *Phys. Fluids* **12**, 1409 (1969) (C.S. Liu).
PPC-99 "Transverse Plasma Echo Phenomena in a Hot Collisionless Plasma", *Phys. Fluids* **12**, 1409 (1969) (C.S. Liu).
PPC-100 "Drift Waves in the Linear Regime", *Phys. Fluids* **13**, 641 (1970) (R. Roehrig and A.Y. Wong).
PPC-101 "Parametric Modulation Coupling Between Drift Waves in Plasmas", *Phys. Fluids* **13**, 672 (1970) (F. Hal and A.Y. Wong).
PPC-102 "Nonlinear Oscillatory Phenomena with Drift Waves in Plasmas", September 1970 (F. Hal and A.Y. Wong).
PPC-103 "Ion-Burst Excited by a Grid in a Plasma", *J. Applied Phys.* **41**, 758 (1970) (H. Ileri and R.J. Taylor).
PPC-104 "Measurements of Diffusion in Velocity Space from Ion-Ion Collisions", *Phys. Rev.* **182**, 1 (1969) (A. Wong and D. Baker).
PPC-105 "Nonlinear Excitation in the Ionosphere", March 1969 (A.Y. Wong).
PPC-106 "The First Order Ion Energy Distribution in Ion Acoustic Waves", *Phys. Rev. Lett.* **22**, 923 (1969) (H. Ileri and R. Taylor).
PPC-107 "A New Representation for the Conductivity Tensor of a Collisionless Plasma in a Magnetic Field", *Festschrift for Gregor Metzger* (U. of Chicago Press, 1969) March 1969 (B.D. Fried and C. Hedrick).
PPC-108 "Direct Measurements of Linear Growth Rates and Nonlinear Saturation Coefficients", *Phys. Rev. Lett.* **23**, 163 (1969) (A.Y. Wong and F. Hal).
PPC-109 "Electron Precipitation Nucleation", *J. Geophys. Res.* **74**, 1279 (1970) (F. Coroniti and C.F. Kennel).
PPC-110 "Auroral Microprecipitation Instability", *J. Geophys. Res.* **75**, 1863 (1970) (F. Coroniti and C.F. Kennel).
PPC-111 "Effect of Magnetic Collisions on the Growth of the Ionospheric E Layer", *Phys. Fluids* **13**, 154 (1970) (A.Y. Wong and F. Hal).
PPC-112 "Theory of Stability of Large Amplitude Periodic (BDS) Waves in Collisionless Plasmas", *Phys. Fluids* **13**, 1281 (1970) (A.Y. Wong and F. Hal).
PPC-113 "Observation of Strong Ion Wave-Mode Interaction", August 1969 (R. Taylor and H. Ileri).
PPC-114 "Optical Mixing in a Magnetoactive Plasma", *Phys. Fluids* **13**, 1802 (1970) (G. Weyl).
PPC-115 "Trapped Particles in a Magnetoactive Plasma", *Phys. Rev. Lett.* **25**, 958 (1969) (A.Y. Wong and R. Taylor).
PPC-116 "Formation and Interaction of Ion-Acoustic Solitons", *Phys. Rev. Lett.* **23**, 11 (1970) (H. Ileri, R.J. Taylor and D.R. Baker).
PPC-117 "Turbulent Loss of Ring Current Protons", *J. Geophys. Res.* **75**, 6699 (1970) (F.V. Coroniti and R.M. Thorne).
PPC-118 "Efficient Modulation Coupling Between Electron and Ion Resonances in Magnetoactive Plasmas", *Phys. Rev. Lett.* **24**, 804 (1970) (A.Y. Wong, D.R. Baker and H. Ileri).
PPC-119 "Interaction of Quasi-Transverse and Quasi-Longitudinal Waves in an Inhomogeneous Vlasov Plasma", Dissertation, January 1970 (C.L. Hedrick).
PPC-120 "Observation of Strong Ion-Acoustic Wave-Mode Interaction", January 1970 (R.J. Taylor and H. Ileri).
PPC-121 "Perturbed Ion Distributions in Ion Waves and Echoes", *Phys. Fluids* **13**, 2348 (1970) (H. Ileri and R.J. Taylor).
PPC-122 "The Analytic and Asymptotic Properties of the Plasma Dispersion Function", February 1970 (A. Mann and G. Johnston).
PPC-123 "Effect of Ion-Ion Collision and Ion Wave Turbulence on the Ion Wave Echo", June 1970 (D. Baker).
PPC-124 "Dispersion Discontinuities of Strong Collisionless Shocks", *J. Plasma Phys.* **4**, 265 (1970) (F.V. Coroniti).
PPC-125 "An Ion Cyclotron Instability", Dissertation, April 1970 (E.S. Weibel).
PPC-126 "Turbulence Structure of Finite Beta Perpendicular Fast Shocks", *J. Geophys. Res.* **75**, 7007 (1970) (F.V. Coroniti).
PPC-127 "Stepping of Ion Acoustic Waves and Formation of Collisionless Electrostatic Shocks", April 1970 (R. Taylor).
PPC-128 "A Method of Studying Trapped Particles Behavior in Magnetic Geometries", *Phys. Rev. Lett.* **25**, 1762 (1970) (C.S. Liu and A.Y. Wong).
PPC-129 "A Note on the Differential Equation $g'' + 2g = 0$ ", April 1970 (E.S. Weibel).
PPC-130 "Plasma Response to a Step Electric Field Greater than the Critical Runaway Field, With and Without an Externally Applied Magnetic Field", Dissertation, June 1970 (J.C. Robinson).
PPC-131 "The Application of an On-Line System as a Tool for Teaching Physics", *Proc. of Computer Graphics*, 136 (1970) (B.D. Fried and R.B. White).
PPC-132 "High Frequency Hall Current Instability", *Radio Sci.* **5**, 209 (1971) (K. Lee, C.F. Kennel and J.M. Kindel).
PPC-133 "Linear Wave Train Structure of Collisionless Magnetic Slow Shocks", *Nuclear Fusion* **11**, 261 (1971) (F.V. Coroniti).
PPC-134 "Field-Aligned Current Instabilities in the Tropic Ionosphere", *J. Geophys. Res.* **76**, 3005 (1971) (J.M. Kindel and C.F. Kennel).
PPC-135 "Spatial Cyclotron Damping", *Phys. Fluids* **15**, 160 (1972) (C. Olson).
PPC-136 "Electromagnetic Plasma Wave Propagation Along a Magnetic Field", Dissertation, September 1970 (C. Olson).
PPC-137 "Radiationless Plasma Wave-Strapping Electron Bursts", *Phys. Fluids* **15**, 1997 (1972) (R. Ileri, P.J. Barrett, R.B. White and A.Y. Wong).
PPC-138 "Relativistic Electron Precipitation During Magnetic Storm Main Phase", *J. Geophys. Res.* **76**, 4446 (1971) (R.M. Thorne and C.F. Kennel).
PPC-139 "A Unified Theory of SAR-Arc Formation at the Plasmapause", *J. Geophys. Res.* **76**, 4428 (1971) (J.M. Cornwall, F.V. Coroniti and R.M. Thorne).
PPC-140 "Nonlinear Collisionless Interaction Between Electron and Ion Modes in Inhomogeneous Magnetoactive Plasmas", Dissertation, December 1970 (M. Booth).
PPC-141 "Oblique Resonance Coupling of Radar Energy to Ionospheric Irregularities", *Comments on Astrophys. and Space Phys.*, **2**, 87 (1971) (C.F. Kennel).
PPC-142 "Current-Driven Electrostatic and Electromagnetic Ion Cyclotron Instabilities", February 1971 (D.M. Forslund, C.F. Kennel and J.M. Kindel).
PPC-143 "Ion Acoustic Waves in a Multi-Ion Plasma", *Phys. Fluids* **14**, 2388 (1971) (B.D. Fried, R.B. White and T. Samc).
PPC-144 "Locating the Magnetospheric Ring Current", *Comments Astrophys. and Space Phys.* **3**, 115 (1971) (C.F. Kennel and R. Thorne).
PPC-145 "Ionospheric Turbulence and the Ring Current", *Phys. Rev. Lett.* **27**, 973 (1971) (A.Y. Wong and R. Thorne).
PPC-146 "Density Step Production of Large Amplitude Collisionless Electrostatic Shocks and Solitons", June 1971 (D.B. Cohen).
PPC-147 "Turbulent Resistivity, Diffusion and Heating", *Plasma Phys. and Controlled Nuclear Fusion Res.* **2**, Vienna (1971) (B.D. Fried, C.F. Kennel, K. McKemie, F.V. Coroniti, J.M. Kindel, R. Stenzel, R.J. Taylor, R.B. White, A.Y. Wong, M. Bernstein, J.M. Sellen, D. Forslund and R.A. Sagdeev).
PPC-148 "Nonlinear Evolution and Saturation of an Unstable Electron Wave", August 1971 (B.D. Fried, C.S. Liu, R.M. Mann, and R.A. Sagdeev).
PPC-149 "Cross-Field Currents and Ion Acoustic Instability", *Phys. Rev. Lett.* **28**, 337 (1972) (P.J. Barrett, B.D. Fried, C.F. Kennel and R.A. Sagdeev).
PPC-150 "3-D Velocity Space Diffusion in Beam-Plasma Interaction Without Magnetic Field", *Proc. on 3rd Int'l. Conf. on Quasilinear Plasmas*, Eilat, 291 (1971) (P.J. Barrett, D. Gressillon and A.Y. Wong).
PPC-151 "Dayside Auroral Oval Plasma Density and Conductivity Enhancements Due to Magnetosheath Electron Precipitation", *J. Geophys. Res.* **77**, 2294 (1972) (C.F. Kennel and M.H. Rees).
PPC-152 "Collisionless Wave-Particle Interactions Perpendicular to the Magnetic Field", *Phys. Rev. Lett.* **29**, 41 (1972) (A.Y. Wong and D.L. Jassby).
PPC-153 "Magnetospheric Substorms", *Cosmic Plasma Physics*, 1971 (Plenum Press, Karl Schindler, 1972) (F.V. Coroniti and C.F. Kennel).
PPC-154 "Magnetopause Motions, DP-2, and the Growth Phase of Magnetospheric Substorms", September 1971 (F.V. Coroniti and C.F. Kennel).
PPC-155 "Structure of Ionospheric Solitons and Shock Waves in a Two-Component Plasma", *Phys. Fluids* **15**, 1484 (1972) (R.B. White, B.D. Fried and F.V. Coroniti).
PPC-156 "Solar Wind Interaction with Lunar Magnetic Field", *J. Plasma Phys.* November 1971 (G. Siscoe and B. Goldstein).
PPC-157 "Changes in Magnetospheric Configuration During Substorm Growth Phase", *J. Geophys. Res.* **77**, 3361 (1972) (F.V. Coroniti and C.F. Kennel).
PPC-158 "Trip Report-1971 Kiev Conference on Plasma Theory and Visits to Lebedev and Kurchatov Institutes", October 1971 (B.D. Fried).
PPC-159 "Pitch-Angle Diffusion of Radiation Belt Electrons Within the Plasmasphere", *J. Geophys. Res.* **77**, 3455 (1972) (L.R. Lyor and R.M. Thorne).
PPC-160 "Stability of a High Beta Plasma", *Phys. Fluids* **15**, 1864 (1972) (F. Chen, D. Jassby and M. Maricic).
PPC-161 "Remote Plasma Control, Heating Measurements of Electron Distribution and Trapped Particles by Nonlinear Electromagnetic Interaction", *J. Plasma Phys. and Controlled Nuclear Fusion Res.* **3**, 335 (1971) (A.Y. Wong, F.F. Chen, N. Booth, D.L. Jassby, R. Stenzel, D. Baker and C.S. Liu).
PPC-162 "Computational and Experimental Plasma Physics for Theoreticalians", *Adv. Plasma Phys.* **5**, 211 (1974) (B.D. Fried).
PPC-163 "Threshold and Saturation of the Parametric Decay Instability", *Phys. Rev. Lett.* **28**, 274 (1972) (R. Stenzel and A.Y. Wong).
PPC-164 "Laser Amplification in an Inhomogeneous Plasma", January 1972 (R.B. White).
PPC-165 "Electron Beam Instability and Control of Electrojet Irregularities", *J. Geophys. Res.* **77**, 4197 (1972) (K. Lee, P.K. Kaw and C.F. Kennel).
PPC-166 "Ion Heating Via Turbulent Ion Acoustic Waves", *Phys. Rev. Lett.* **29**, 34 (1972) (R.J. Taylor and F.V. Coroniti).
PPC-167 "Polarization of the Auroral Electrojet", *J. Geophys. Res.* **77**, 2635 (1972) (F.V. Coroniti and C.F. Kennel).
PPC-168 "Mode Coupling and Wave Particle Interactions for Unstable Ion Acoustic Waves", *Phys. Fluids* **15**, 2275 (1972) (F. Martin and B.D. Fried).
PPC-169 "Parallel Magnetic Multi-Pole Confinement of a Magnetic Field-Free Plasma", Dissertation, March 1972 (R. Limpaecher).
PPC-170 "Unstable Ionospheric Shock Waves", Dissertation, April 1972 (R.M. Mann).
PPC-171 "Auroral Plasma Instability in a Magnetoactive Plasma", Dissertation, April 1972 (C. Olson).
PPC-172 "Radiationless Plasma Wave-Strapping Electron Bursts", *Phys. Fluids* **15**, 1997 (1972) (R. Ileri, P.J. Barrett, R.B. White and A.Y. Wong).
PPC-173 "Radiationless Plasma Wave-Strapping Electron Bursts", *Phys. Fluids* **15**, 1997 (1972) (R. Ileri, P.J. Barrett, R.B. White and A.Y. Wong).
PPC-174 "A New Role for Infrared Lasers", *Comments on Plasma Phys. and Cont. Fusion* **3**, 81 (1973) (F. Chen).
PPC-175 "Electrostatic Instability of Ring Current Protons Beyond the Plasmapause During Injection Events", *J. Geophys. Res.* **77** (1972) (F. Chen).
PPC-176 "Magnetospheres of the Outer Planets", *Space Sci. Rev.* **13**, 511 (1973) (C.F. Kennel).
PPC-177 "Measurement of Transverse and Longitudinal Heat Flow in a Laser-Heated, Magnetically Confined Arc Plasma", Dissertation, June 1972 (S.M. Fay).
PPC-178 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-179 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-180 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-181 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-182 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-183 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-184 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-185 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-186 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-187 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-188 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-189 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-190 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-191 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-192 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-193 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-194 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-195 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-196 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-197 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-198 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-199 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).
PPC-200 "Plasma Physics of the Earth's Ionosphere", *Phys. Rev. Lett.* **26** (1972) (S.M. Fay, R. Stenzel, R.M. Thorne and R.E. Holzer).

- PHC-124 "Calculation of Reflection and Transmission Coefficients for a Class of One-Dimensional Wave Propagation Problems in Inhomogeneous Media," *Phys. Rev. Lett.*, **35**, 1565 (1975) (A. B. Fiedel and J. P. Fiedel).
- PHC-125 "Inhomogeneous Media Functions for Parallel Plasma Density Profiles," *J. Plasma Phys.*, **15**, 275 (1975) (A. B. Fiedel and R. B. White).
- PHC-126 "Calculation of Electromagnetic Waves in Overdense Plasma," *J. Plasma Phys.*, **15**, 585 (1975) (A. B. Fiedel and R. B. White).
- PHC-127 "Abstracts presented at the Plasma Physics Annual Meeting, November 13-16 (1975)."
- PHC-128 "Can the Ionosphere Regulate Magnetospheric Convection?" *J. Geophys. Res.*, **78**, 7837 (1973) (F. V. Coroniti and C. F. Kennel).
- PHC-129 "Nonlinear Landau Damping of Oscillating Two-Stream Instability," *Phys. Fluids*, **16**, 1580 (1973) (K. Nishikawa, Y. C. Lee and P. K. Kaw).
- PHC-130 "Finite Beta Plasma," Dissertation, October (1972) (M. S. Chance).
- PHC-131 "Wave Packet Formation of Nonlinear Plasma Waves," *Phys. Fluids*, **16**, 1321 (1973) (K. Nishikawa and B. D. Fried).
- PHC-132 "Nonlinear Cyclotron Drift Instability," *Phys. Fluids*, **16**, 1321 (1973) (K. Nishikawa and B. D. Fried).
- PHC-133 "Resonant Excitation of Electromagnetic Waves in Plasma," *Phys. Fluids*, **16**, 1676 (1973) (G. Schmidt).
- PHC-134 "Resonant Excitation of Electromagnetic Waves in Plasma," *Phys. Fluids*, **16**, 1676 (1973) (G. Schmidt).
- PHC-135 "Energy Ion Beam Source and Free-Stream Beam Diagnostic Techniques," *Rev. Sci. Instrum.*, **44**, 617 (1973) (R. L. Stenzel and B. H. Rippl).
- PHC-136 "Electron Plasma Waves in an Unbounded Uniform Magnetoplasma," *Phys. Fluids*, **16**, 565 (1973) (R. L. Stenzel).
- PHC-137 "Theoretical Calculation of Type I Irregularities in the Equatorial Electrojet," *J. Geophys. Res.*, **78**, 4619 (1973) (K. Lee and C. F. Kennel).
- PHC-138 "Effects of Propagation Parallel to the Magnetic Field on the Type I Electrojet Irregularity Instability," *Planetary and Space Sci.*, **21**, 1339 (1973) (K. Lee and C. F. Kennel).
- PHC-139 "Modeling Simulation of Parametric Instability," November (1972) (R. L. Stenzel).
- PHC-140 "Theory of Double Resonance Parametric Excitation in Plasmas," *Phys. Fluids*, **16**, 2270 (1973) (D. Arnush, B. D. Fried, C. F. Kennel, K. Nishikawa and A. Y. Wong).
- PHC-141 "Finite Beta Drift Angle Instability," *J. Geophys. Res.*, **78**, 7521 (1973) (M. S. Chance, F. V. Coroniti and C. F. Kennel).
- PHC-142 "The Formation of Ion Acoustic Shocks," *Phys. Fluids*, **17**, 217 (1974) (R. B. White, B. D. Fried and F. V. Coroniti).
- PHC-143 "Experiments on Parametric Instabilities," March (1973) (A. Y. Wong).
- PHC-144 "On Cosmic Ray Generation by Pulsars," *Phys. Rev. Lett.*, **31**, 1364 (1973) (C. F. Kennel, G. Schmidt and T. Wilcox).
- PHC-145 "On the Propagability of Stable Stationary Spectra of Unstable Type I Equatorial Electrojet Irregularities," *J. Geophys. Res.*, **78**, 7837 (1973) (F. V. Coroniti and C. F. Kennel).
- PHC-146 "Special Group Properties of Parametric and Backscattering Plasma Instabilities," April (1973) (B. D. Fried, R. Gould and G. Schmidt).
- PHC-147 "Evolution of K_z-Like Modes with Trapped Electrons," *Phys. Rev. Lett.*, **30**, 1299 (1973) (A. Y. Wong, B. H. Quon and B. Rippl).
- PHC-148 "Stabilization of Ion Acoustic Waves by Electron Trapping," *Phys. Fluids*, **17**, 206 (1974) (N. Albricht).
- PHC-149 "Turbulence in Electromagnetic Ion Acoustic Shocks," *Phys. Fluids*, **16**, 2304 (1973) (R. M. Means, F. V. Coroniti, A. Y. Wong and B. H. Quon).
- PHC-150 "Thermal Ionization Function in a Magnetized Plasma," June (1973) (Y. C. Lee and C. S. Liu).
- PHC-151 "Physical Interpretation of the Oscillatory Two-Stream Instability," June (1973) (A. Y. Wong and G. Schmidt).
- PHC-152 "Relativistic Particle Motion in Nonuniform Electromagnetic Waves," *Phys. Rev. Lett.*, **31**, 1380 (1973) (G. Schmidt and T. Wilcox).
- PHC-153 "The Ring Current and Magnetic Storms," *Radio Sci.*, **8**, 1007 (1973) (F. V. Coroniti).
- PHC-154 "Energetic Electrons in Jupiter's Magnetosphere," *Astrophys. J.*, **27**, 261 (1974) (F. V. Coroniti).
- PHC-155 "Stably Trapped Proton Flares in the Jovian Magnetosphere," *Astrophys. J.*, **189**, 383 (1974) (F. V. Coroniti, C. F. Kennel and R. M. Thorne).
- PHC-156 "Nonlinear Scattering Instabilities in an Inhomogeneous Plasma," *Phys. Rev. Lett.*, **31**, 1197 (1973) (J. F. Drake and Y. C. Lee).
- PHC-157 "Growth and Saturation of the Absolute Electron Cyclotron Drift Instability," *Phys. Rev. Lett.*, **31**, 1545 (1973) (B. L. Stenzel and B. H. Rippl).
- PHC-158 "Parametric Instabilities of Electromagnetic Waves in Plasmas," *Phys. Fluids*, **17**, 718 (1974) (J. Drake, P. K. Kaw, Y. C. Lee, G. Schmidt, C. S. Liu and H. M. Rombach).
- PHC-159 "Nonlinear Spectra of Plasmas," Survey Lecture of Int'l. Congress on Waves and Instabilities in Plasmas, Imbroschi, Austr (1973) (G. Schmidt, C. S. Liu and H. M. Rombach).
- PHC-160 "Theoretical Studies of Nonlinear Interactions in Plasmas," *Phys. Fluids*, **17**, 291 (1974) (F. V. Coroniti).
- PHC-161 "Trip Report - 6th European Conference on Cont. Fusion and Plasma Phys., July 30-August 4, 1973," August (1973) (B. D. Fried).
- PHC-162 "Abstracts presented at Philadelphia Meeting of Am. Phys. Soc. Div. Plasma Phys., October 31-November 3, 1973."
- PHC-163 "Enhancement of Plasma DC Currents by Heating AC Fields," *Phys. Fluids*, **17**, 987 (1974) (A. T. Lin and J. M. Dawson).
- PHC-164 "Temporal Electrostatic Instabilities in Inhomogeneous Plasmas," *Phys. Rev. Lett.*, **32**, 135 (1974) (G. Lee and P. K. Kaw).
- PHC-165 "Y. C. Lee and B. H. Rippl," *Phys. Rev. Lett.*, **32**, 457 (1974) (G. Lee and B. H. Rippl).
- PHC-166 "Backscattering Decay Processes in Electron Beam-Plasma Interactions Including Ion Dynamics," *Phys. Rev. Lett.*, **32**, 406 (1974) (B. H. Quon, A. Y. Wong and B. H. Rippl).
- PHC-167 "Conversion of Electromagnetic Waves to Electrostatic Waves in Inhomogeneous Plasmas," *Phys. Rev. Lett.*, **32**, 654 (1974) (R. Stenzel, A. Y. Wong and B. H. Quon).
- PHC-168 "Langmuir Turbulence - Condensation and Collapse," *Com. Plasma Phys. and Cont. Fusion* (1974) (Y. C. Lee, C. S. Liu and R. Stenzel).
- PHC-169 "The Consequences of Microturbulence on Geomagnetically Trapped Particles," *Rev. Space Sci.*, **16**, 443 (1974) (R. M. Thorne).
- PHC-170 "Linear Wave Conversion in Inhomogeneous Plasmas," March (1974) (D. L. Kelly and A. Kono).
- PHC-171 "The Cause of Storm After-effects in the Middle Latitude D-region Ionosphere," *J. Atmospheric and Terrestrial Phys.* (1974).
- PHC-172 "Application of an Electromagnetic Particle Simulation Code to the Generation of Electromagnetic Radiation," *Phys. Fluids*, **17**, 1995 (1974) (A. T. Lin, J. M. Dawson and H. Ombak).
- PHC-173 "The 1965 Lunar Flare in a Magnetized Plasma," *Phys. Fluids*, **17**, 1995 (1974) (A. T. Lin, J. M. Dawson and H. Ombak).
- PHC-174 "What we have Learned from the Study of a Magnetized Plasma," *Phys. Fluids*, **17**, 1995 (1974) (A. T. Lin, J. M. Dawson and H. Ombak).
- PHC-175 "Observation of the Parametric Instability and Oscillating Two-Stream Instability," April (1974) (H. C. Kiu, R. Stenzel and A. Y. Wong).
- PHC-176 "Linear Beam Plasma Interaction Including Ion Dynamic," Dissertation, June (1974) (R. H. Quon).
- PHC-177 "Linear Conversion and Parametric Instabilities in a Non-Uniform Plasma," *Phys. Fluids*, **17**, 1995 (1974) (R. H. Quon).
- PHC-178 "Quasilinear Scattering of Electromagnetic Waves in Plasmas," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-179 "The Effect of the Parametric Instability on the Propagation of a Wave in a Magnetized Plasma," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-180 "Effect of the Parametric Instability on the Propagation of a Wave in a Magnetized Plasma," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-181 "Quasilinear Scattering of Electromagnetic Waves in Plasmas," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-182 "Response of the Ionosphere to a Geomagnetic Storm," *J. Atmos. and Terr. Phys.* (1974) (J. P. Schieldge and S. V. Venkateswaran).
- PHC-183 "The Effect of the Parametric Instability on the Propagation of a Wave in a Magnetized Plasma," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-184 "The Effect of the Parametric Instability on the Propagation of a Wave in a Magnetized Plasma," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-185 "The Effect of the Parametric Instability on the Propagation of a Wave in a Magnetized Plasma," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-186 "The Effect of the Parametric Instability on the Propagation of a Wave in a Magnetized Plasma," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-187 "The Effect of the Parametric Instability on the Propagation of a Wave in a Magnetized Plasma," *Phys. Fluids*, **16**, 201 (1973) (A. T. Lin and J. M. Dawson).
- PHC-188 "The Effect of the Param

- PPG-246 "Parametric Instabilities with Finite Wavelength Pump", B.D. Fried, T. Ikemura, K. Nishikawa, and G. Schmidt, *Phys. of Fluids*, 19, 1975 (1976).
- PPG-247 "The Scattering of Cosmic Rays by Magnetic Bubbles", R.F. Flewelling & F.V. Coroniti, *Astrophys. J.*, 205, 135 (1976).
- PPG-248 "Plasma Simulation on the CHI Microprocessor System", T. Kamimura, J.M. Dawson, B. Rosen, G.J. Culler, R.D. Levee and G. Ball., (1975).
- PPG-249 "Nonlinear Interactions of Focused Resonance Cone Fields with Plasmas", R. Stenzel and W. Geikelman, *Phys. Fluids* 20, 108 (1977).
- PPG-250 "The Effect of Pump Cutoff on Parametric Instabilities in an Inhomogeneous Plasma", J. VanDam and Y.C. Lee, (1976).
- PPG-251 "Parametric Excitation of Ion Density Fluctuations in the Relativistic Beam-Plasma Interaction", H. Schamel, Y.C. Lee, & G.J. Morales, *Phys. Fluids* 19, 849 (1976).
- PPG-252 "The Spiky Turbulence Generated by a Propagating Electrostatic Wave of Finite Spatial Extent", G.J. Morales, & Y.C. Lee, *Phys. Fluids* 19, 690 (1976).
- PPG-253 "Departures from Theory of the Experimental Line Profile of Helium I 4471.5A", J. Turechek, January (1976).
- PPG-254 "Maintenance of the Middle Latitude Nocturnal D-Layer by Energetic Electron Precipitation", W. Spjeldvik & R. Thorne, *Pure and Applied Geophys.* 114, 497 (1976).
- PPG-255 "Free Boundaries for Plasmas in Surface Magnetic Field Configurations", B.D. Fried, J.W. VanDam & Y.C. Lee, *Phys. of Fluids*, 19, 1909 (1976).
- PPG-256 "Soliton-Like Structures in Plasmas", Y.C. Lee and G.J. Morales, March (1976). To appear in *Rocky Mtn. Math. J.*
- PPG-257 "Magnetic Field Reconnection in a Collisionless Plasma", F.V. Coroniti and A. Eviatar, *Astrophys. J. Supp.* 33, 189 (1977).
- PPG-258 "The Earth's Magnetosphere", F.V. Coroniti, Accepted by *Ency. of Phys.* (1976).
- PPG-259 "Lossy Radial Diffusion of Relativistic Jovian Electrons", D.D. Barbosa and F.V. Coroniti, *J. Geophys. Res.* 81, 4553 (1976).
- PPG-260 "Relativistic Electrons and Whistlers in Jupiter's Magnetosphere", D.D. Barbosa and F.V. Coroniti, *J. Geophys. Res.* 81, 4531 (1976).
- PPG-261 "A Self-Consistent Magnetostatic Particle Code for Numerical Simulation of Plasmas", J. Busnardo-Neto, P.L. Pritchett, A.T. Lin & J.M. Dawson, *J. Comp. Phys.* 23, 300 (1977).
- PPG-262 "Relativistic Effects in Resonance Absorption", J.F. Drake and Y.C. Lee, *Phys. of Fluids* 19, 1772 (1976).
- PPG-263 "Convective Electron Loss Cone Instabilities", M. Ashour-Abdalla and C.F. Kennel. Submitted to *J. Geophys. Res.* (1976).
- PPG-264 "Experiments on Parametric Instabilities in Laser-Beam Plasma Interactions", R. Chen, Presented at Nobel Symposium.
- PPG-265 "The Local Time Variation of ELF Emissions During Periods of Substorm Activity", R.M. Thorne, F.R. Church, W.J. Malloy, B.T. Tsurutani, *J. of Geophys. Res.*, 82, 1585 (1977).
- PPG-266 "An Investigation of Relativistic Electron Precipitation Events and their Association with Magnetospheric Substorm Activity", R.M. Thorne and T.R. Larsen, *J. of Geophys. Res.* 81, 5501 (1976).
- PPG-267 "The Free Electron Laser", T. Kwan, J.M. Dawson & A.T. Lin, *Phys. Fluids* 20, 581 (1977).
- PPG-268 "Stimulated Compton Scattering of Electromagnetic Waves off Plasma Ions", A.T. Lin and J.M. Dawson, *Phys. Fluids* 20, 538 (1977).
- PPG-269 "Stability of Shell Distributions", D.D. Barbosa, Submitted to *Planetary and Space Sciences*, July (1976).
- PPG-270 "Synchro-Compton Radiation Damping of Relativistically Strong Plasma Waves", C.F. Kennel, E. Asseo and R. Pellat, Submitted to *Astronomy and Astrophysics*, July (1976).
- PPG-271 "Particle and Wave Dynamics in a Magnetized Plasma Subject to High RF Pressure", W. Geikelman & R.L. Stenzel, *Phys. Fluids* 20, 1316 (1977).
- PPG-272 "Energetic Radiation Belt Electron Precipitation; A Natural Depletion Mechanism for Stratospheric Ozone, R.M. Thorne, *Science*, 195, 287 (1977).
- PPG-273 "CTR Using the P-1^B Reaction", J.M. Dawson, August (1976).
- PPG-274 "San Francisco Abstracts - Papers to be Presented at San Francisco Meeting of the American Physical Society Division of Plasma Physics, November 15-19 (1976).

- PPG-275 "Surface Multipole Guide Field for Plasma Injection", R.A. Breun, B.H. Rael, & A.Y. Wong, Review of Scientific Instruments 48 (1977).
- PPG-276 "Higher Order Adiabatic Invariants and Non-Adiabatic Diffusion for Geometric Mirrors", Y.C. Lee, T.K. Samec, & B.D. Fried, Phys. Fluids, 20, 815 (1977).
- PPG-277 "Electromagnetic Interactions with Inhomogeneous Plasmas", A.Y. Wong. Submitted to Nuclear Fusion and RPI Oct. (1976).
- PPG-278 "The Importance of Electrostatic Ion-Cyclotron Instability for Quiet-Time Proton Auroral Precipitation", M. Ashour-Abdalla & R.M. Thorne, Geophys. Res. Lett., 4, 45, (1977).
- PPG-279 "Self Channeling of Whistler Waves", E.S. Weibel, Phys. Letts. 61, 37 (1977).
- PPG-280 "Formation of Potential Double Layers in Plasmas", B.H. Quon & A.Y. Wong, Phys. Rev. Letts., 37, 1393 (1976).
- PPG-281 "Plasma Confinement by Surface Magnetic Fields", T.K. Samec.
- PPG-282 "Kinetic Theory of Tearing Instabilities", J.F. Drake & Y.C. Lee, Phys. Fluids 20, 1341 (1977).
- PPG-283 "Jupiter's Magnetosphere", C.F. Kennel & F.V. Coroniti, Annual Rev. of Astro. & Astrophys., Submitted Nov., 1976.
- PPG-284 "Direct Density Display with a Resonance Cone RF Probe", J. Ickovic, R.L. Stenzel & W. Gekelman, Rev. of Scientific Instr. 48, 435 (1977).
- PPG-285 "The Ponderomotive Effect in a Magnetized Plasma", E.S. Weibel.
- PPG-286 "The Fluid Theory of the Ponderomotive Effect in a Magnetized Plasma", E.S. Weibel.
- PPG-287 "Possible Origins of Time Variability in Jupiter's Outer Magnetosphere", F.V. Coroniti & C.F. Kennel, Geophys. Res. Lett., 4, 287 (1977).
- PPG-288 "Coupling of Lower-Hybrid Radiation at the Plasma Edge", G.J. Morales, Phys. Fluids, 20, 1164 (1977).
- PPG-289 "Nonlinear Evolution of Collisionless and Semi-Collisional Tearing Modes", J.F. Drake and Y.C. Lee, Phys. Rev. Lett. 39, 453 (1977).
- PPG-290 "Model Analysis for Shear Stabilization of Collisionless Drift Waves", Y.C. Lee & T. Tange.
- PPG-291 "Particle Orbits and Loss Regions in UCLA Tokamaks", C.P. Lee, January (1977)
- PPG-292 "Generation of Density Cavities and Localized Electric Fields in a Nonuniform Plasma", G.J. Morales & Y.C. Lee, Phys. Fluids, 20, 1135 (1977).
- PPG-293 "Simulation Study of Bernstein Modes", T. Kamimura, T. Wagner & J.M. Dawson. Submitted to Phys. Fluids (1977).
- PPG-294 "Surface Trappings of Oxygen in Tokamaks and its Removal", Lena Oren and R.J. Taylor.
- PPG-295 "Low-Z Impurity Control for Reactor Size Tokamaks", R.J. Taylor and L. Oren.
- PPG-296 "Computer Simulation of Current Penetration in a Plasma", A.T. Lin and J.M. Dawson, Submitted to Phys. Fluids, March, 1977.
- PPG-297 "The Alfvén-Ion Cyclotron Instability: Its Physical Mechanism and Observation in Computer Simulation", T. Tajima, K. Mima, and J.M. Dawson, Phys. Rev. Lett. 39, 201 (1977).
- PPG-298 "Phase Mixing in the Continuous Spectrum of Alfvén Waves", P.L. Pritchett & J.M. Dawson, Phys. Fluids, Submitted May, (1977).
- PPG-299 "Nonlinear Evolution of the Collisionless Tearing Mode", F.V. Coroniti, Phys. Rev. Lett. 38, 1355 (1977).
- PPG-300 "Active Stimulation of Ionospheric Plasma", A.Y. Wong, April, 1977.
- PPG-301 "Confinement of High β Toroidal Plasmas and Currents in Surface Magnetic Fields", D. Mamas, R.W. Schumacher, A.Y. Wong and R.A. Breun, submitted to Phys. Rev. Lett.
- PPG-302 "Super Conducting Surmacks", A. Wong and L. Miller, May (1977).
- PPG-303 "Plasma Heating by a Relativistic Electron Beam with Secondary Instabilities", T. Tajima. Submitted to J. Plasma Phys.
- PPG-304 "Langmuir Waves in a Spatially Periodic Potential", A. Banos, E.S. Weibel, and J.W. VanDam, Phys. Fluids, submitted June, 1977.
- PPG-305 "Quasi-Linear Theory for Non-Uniform Turbulence", E.S. Weibel, June, 1977.

- PPG-306 "Kinetic Theory of the Internal $m = 1$ Resistive Mode," J. F. Drake, Y. C. Lee, L. Chen, P. H. Rutherford, P. K. Kaw, J. Y. Hsu, and C. S. Liu. Submitted to Phys. Fluids.
- PPG-307 "Some Safety Considerations in Laser Controlled Thermonuclear Reactors," T. E. Botts, D. Breton, C. K. Chan, S. I. Levy, M. Sehner, and A. Z. Ullman.
- PPG-308 "Some Safety Considerations for Conceptual Tokamak Fusion Power Reactors," W. E. Kastenber, and D. Okrent.
- PPG-309 "Coupling of Tearing Modes by Toroidal Effects," J. F. Drake. July, 1977.
- PPG-310 "Stabilization of the Alfvén-Ion Cyclotron Instability in Inhomogeneous Media," T. Tajima and K. Mima, submitted to Phys. Fluids.
- PPG-311 "Particle-Caviton Interactions," A. Y. Wong, P. Leung, and D. Eggleston. Submitted to Phys. Rev. Lett., August, 1977.
- PPG-312 "Nonlinear Evolution of Drift Alfvén and Double Tearing Instabilities," A. T. Lin. Submitted to Phys. Fluids, August, 1977.
- PPG-313 "Atlanta Abstracts - Papers to be Presented at Atlanta Meeting of the American Physical Society Division of Plasma Physics, November 7 - 11, 1977."
- PPG-314 "Self-Consistent Potential for a Relativistic Magnetized Electron Beam Through a Metallic Boundary," T. Tajima and F. Mako.
- PPG-315 "A Magnetohydrodynamic Particle Code for Fluid Simulation of Plasmas," J. N. Leboeuf, T. Tajima and J. M. Dawson.
- PPG-316 "Stability Theory of a Relativistic Electron Beam-Plasma System with Finite Geometries," T. Tajima.
- PPG-317 "Anomalous Diffusion and Resistivity of a Turbulent Plasma," R. L. Stenzel and W. Gekelman, submitted to Phys. Rev. Lett., September, 1977.
- PPG-318 "Properties of Large Amplitude Langmuir Solitons," M. D'Evelyn and G. J. Morales, submitted to Phys. Fluids, August, 1977.
- PPG-319 "End 'Plugged Long Straight Systems for Thermonuclear Reactors," J. M. Dawson, October, 1977.
- PPG-320 "Diffuse Auroral Precipitation," M. Ashour-Abdalla and C. F. Kennel, October, 1977.
- PPG-321 "Modulational Instability of Strongly Dispersive Drift Waves and Formation of Convective Waves," K. Mima and Y. C. Lee. Submitted to Phys. Rev. Lett., October, 1977.
- PPG-322 "Some Safety Considerations for Conceptual Fusion-Fission Hybrid Reactors," W. Kastenber and D. Okrent. October, 1977.
- PPG-323 "Space-Time Resolved Line Radiation in Microtor-Macrotr Tokamaks," L. Oren and R. J. Taylor.
- PPG-324 "Efficiency of Antenna Coupling and Propagation of Fast Alfvén Modes in Macrotr," G. J. Morales and R. J. Taylor.
- PPG-325 "Topology of Cr Influx in Tokamaks," L. Oren and R. J. Taylor.
- PPG-326 "Introduction to Experimental Plasma Physics, Vol. I," A. Wong, November, 1977.
- PPG-327 "Inductive Reconnection at an X-type Neutral Line," F. V. Coroniti and M. Ashour-Abdalla, November, 1977.
- PPG-328 "Computer Simulation Study of Waves in Bounded Plasma with Application to Lower Hybrid Heating," Viktor Decyk, Dissertation, November, 1977.
- PPG-329 "Double-Layer Forward Shocks in an MHD Fluid," T. Tajima, J. N. Leboeuf, and J. M. Dawson. To be submitted to Phys. Rev. Lett., December, 1977.
- PPG-330 "Particle Orbits and Loss Regions in the UCLA Tokamaks," A. Baños Jr., and C. P. Lee.
- PPG-331 "Computer Model for Bounded Plasma," Viktor K. Decyk and John M. Dawson, submitted to the J. of Computational Physics, December, 1977.
- PPG-332 "Interchange Instabilities in a Compressible Plasma," P. Pritchett, C.C. Wu and J. Dawson. Submitted to Phys. Fluids, December, 1977.
- PPG-333 "CO₂ Laser Thomson Scattering from a Pulsed Hydrogen Arc." W.A. Peebles and M.J. Herbst. January, 1978.
- PPG-334 "Plasma Simulation on the UCLA CHI Computer System," J.M. Dawson, R.H. Huff and C-C Wu. To be published in the Proceedings of the 1978 National Computer Conference. January, 1978.

- PPG-335 "Kinetic Theory of $m = 1$ Internal Instabilities," J.F. Drake, submitted to Phys. Fluids, January, 1978.
- PPG-336 "Magnetic Field Generation by the Rayleigh-Taylor Instability," T. Tajima, K. Mima, and J.N. Leboeuf. To be published in the Proceedings of Topical Meeting of Inertial Confinement Fusion (CLEOS). February, 1978.
- PPG-337 "Kinetic Theory of Ballooning Instabilities," Y.C. Lee and J.W. VanDam. To be published in the Proceedings of the High-Beta Tokamak Theory Workshop, Varenna, 1977 (by the U.S. DOE). February, 1978.
- PPG-338 "Excitation of Lower Hybrid Waves in a Finite Plasma," V.K. Decyk, J.M. Dawson, and G.J. Morales, submitted to Phys. Fluids. February, 1978.
- PPG-339 "Explosive Tearing Mode Reconnection in the Magnetospheric Tail," A.A. Galeev, F.V. Coroniti, and M. Ashour-Abdalla, March (1978).
- PPG-340 "Small Amplitude Waves in a Hot Relativistic Two Fluid Plasma," S. Hyun and C.F. Kennel, March (1978).
- PPG-341 "Nonlinear Evolution of Tearing Instabilities: Violations of Constant ψ ," J.F. Drake, P.L. Pritchett, and Y.C. Lee. Submitted to Phys. Rev. Letts., March (1978).

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) EXCITATION OF LOWER HYBRID WAVES IN A FINITE PLASMA		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER PPG-338V
7. AUTHOR(s) V.K. Decyk, J.M. Dawson and G. J. Morales		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0476 P00004
9. PERFORMING ORGANIZATION NAME AND ADDRESS Physics Department University of California Los Angeles, California 90024		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217		12. REPORT DATE February, 1978
		13. NUMBER OF PAGES 41
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) RF Heating Electromagnetic Waves in Plasmas		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A computer simulation study of the launching of lower hybrid waves by external sources and their propagation in a finite but homogeneous plasma slab is presented. The study makes use of a 2-1/2 dimensional, electrostatic particle code developed for simulation of bounded plasma. A number of antenna configurations are considered: a point source, two finite-length capacitor plates, and a phased array of capacitor plates. When the oscillating frequency does not match a bounded plasma resonance, one can observe		

20. Continued.

resonance cones, energy absorption at the plasma surface, ion cyclotron modulation of the source, and energetic ions, depending on the parameters chosen. The excitation of a bounded plasma resonance is also considered. The nonlinear evolution of the resonance shows that wave-particle interactions and ponderomotive force effects play an important role. The possibility of controlling the electron temperature profile is discussed.